

An Clinicopathological And Survival Analysis Of Non-Squamous Cervical Cancers At AHPGIC

Research Article

B.L.Nayak¹, A.K.Padhy², J. Jmohapatra², M.R.Mohapatra², J. Parija, RDAS², S. Mohanty², S.S. Pattnaik^{2*}, S.K.Giri², N. Panda², S.N.Senapathi², L. Pattnaik², L. Sarangi², S. Padhi², S. Samantray², S. Panda², N. Rout², T. Kar², L. Das², Duttaswar Hota², P.C.Mohapatra², P. Devi²

¹ Professor, Department of Gynae Oncology, AHPGIC, Cuttack.

² Department of Gynaecology, Trained in Gynaecology, SCB Medical College & Hospital, Cuttack.

Abstract

AIM AND OBJECTIVE: An Clinicopathological and survival analysis 22 cases of nonsquamous carcinomas of cervix from 2010- 2020.

Primary objective is to analyse the extent of microscopic parametrial involvement in early invasive non squamous carcinoma with respect to age, size of lesion, depth of invasion.

To analyse the nodal status in clinically early invasive nonsquamous carcinoma with respect to other variable i.e age, tumor size, depth of invasion, parametrial involvement.

SECONDARY OBJECTIVE: was to analyse the survival status, of after radical hysterectomy and adjuvant therapy.

MATERIAL AND METHODS: The hps confirmed cases of nonsquamous carcinoma of cervix were included in the study and few cases were ih confirmed.

INCLUSION CRITERIA: Primary non squamous carcinoma of cervix hps confirmed.

EXCLUSION CRITERIA: Squamous carcinoma and metastatic carcinoma of cervix

METHODS: Regression analysis chi square, log rank test. Kaplan Meyers curve was used for survival analysis.

Results: On regression analysis with dependent variable as the pelvic node was analysed with age, depth of invasion and size of lesion and parametrium, none of factors were significantly influencing the nodal status.

Similarly the parametrium positivity taken as the dependent variable was analysed with age, size of lesion, depth of invasion. none of the factors could predict the parametrial involvement. Overall 5-year survival probability for the patients is 0.770 i.e 55 months with 95% CI (0.570, 1.000). Here, the 5-year survival probability for the patients with parametrium positive status is 0.917 i.e 48 months 95% CI (0.773, 1.000). The log rank test for change in survival between two parametrium groups i.e., negative and positive patients gives chi-square value 3.59 (d.f.=1, p-value = 0.06), which indicates two groups do not differ in their survival.

Disease free survival of the parametrium -ve/ parametrium +ve (60/24 mths). The log rank test for change in survival between two parametrium groups i.e., negative and positive patients gives chi-square value 2.8 (d.f.=1, p-value = 0.1), which indicates two groups do not differ in their disease free survival.

Introduction

The global incidence of cervical cancer burden is disproportionately high in low and middle income countries, where 83% of all new cases and 855 of cervical cancer death occur [1]. India accounts for nearly one fourth of the world's cervical cancer deaths, with 60,078 death and 96, 922 new cases in 2018 [2, 3]. Cervical cancer is the most probe cause of maternal mortality in Indian women.

Adenocarcinoma represent 20-25% of cervical cancers in the industrialized countries.

Most of this is due to relative decrease in incidence of squamous cell carcinoma. In contrast to squamous carcinoma smoking does not increase the risk adenocarcinoma. Squamous and non squamous differ in hpv status [4]. Hpv 18 accounts for 50% of non-squamous cell cancer (adenocarcinoma) of 15% of squamous cell

*Corresponding Author:

S.S Pattnaik, M.B.B.S.(SCBMCH),
Department of Gynaecology, Trained in Gynaecology, SCB Medical College & Hospital, Cuttack.
Tel: 8328953390
E-mail: drsmrutisudhapattnaik@gmail.com

Received: April 13, 2022

Accepted: May 31, 2022

Published: June 13, 2022

Citation: B.L.Nayak, A.K.Padhy, J. Jmohapatra, M.R.Mohapatra, J. Parija, RDAS, S. Mohanty, S.S. Pattnaik, et al., An Clinicopathological And Survival Analysis Of Non-Squamous Cervical Cancers At AHPGIC. *Int J Cancer Stud Res.* 2022;9(1e):149-155.

Copyright: S.S Pattnaik©2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

carcinomas.

Management of adenocarcinoma is same as squamous.

Whee as squamous disseminate via lymphatics and adenocarcinoma haematogenous route [5].

This is evident as as after lymphatic dissemination adenocarcinoma has poor prognosis compared to squamous cell carcinoma.[6] Other evidence in support of haematogenous spread [1]. the largest series of surgically teated cervical cancers demonstrated a significantly highe rate of ovaian metastasis with adenocarcinoma (5%vs .8%p<.01).

2. a study of 367 pts of adenocarcinoma from M.D ANDERSON hospital reported a higher rate of distant metastasis for stage II (46%VS13%) and stage III disease (38%ovs21%) when compared squamous carcinoma.

Thee are are very few studies regading the prognostic factors ie clinical and pathological factors influencing the parametrium positivity and nodal status of non squamous carcinoma. that is the reason they ae ovetreated by multimodality i.e(radical sugery with adjuvant).

Studies done by M.d Anderson on 29 patients of adenosquamous and 97 pts of adenocarcinoma in stage 1b1., with radical hysterectomy. On follow up time to recurrence (7.9mths vs 19 monthsp-.01).

A STUDY OF 163 adenocarcinoma and adenosquamous carcinoma with stage 1A2 to IIB disease treated by radical hysterectomy with o without adjuvant radiation found no difference in ecurrence ate or patterns of ecurrences between the two groups, in both low risk, intermediate risk high isk group.[7]

One hundred patients met the inclusion criteria.

The median age was 35 years (range 22-65), and 51% (51/100) had pure high-grade neuroendocrine carcinoma.

No patient had a tumor > 4 cm or suspected parametrial or nodal disease before surgery. Ten patients (10%) had microscopic parametrial compromise in the final surgical specimens. Ninety-four (94%) patients underwent nodal assessment, and 19 (19%) had positive nodes. Ten patients underwent both sentinel lymph node biopsy and pelvic lymphadenectomy, and none had false-negative findings. Patients with parametrial compromise were more likely to have positive pelvic nodes (80% vs 12%, p<0.0001), and a positive vaginal margin (20% vs 1%, p=0.03). All patients with parametrial compromise had lymphovascular space invasion (100% vs 73%, p=0.10). of the 100 patients, 95 (95%) were recommended adjuvant therapy and 89 (89%) were known to have received it. adjuvant pelvic radiotherapy reduced the likelihood of local recurrence by 62%.[8] gloria salvo et al.

Descriptive Statistics

A total 22 cases of non squamous carcinoma were taken for statistical evaluation The median age incidence is 46 yrs. The youngest age is 28 yrs and maximum age 62 yrs. 9(40.9) were <46 yrs and noscaese more than 46 yrs 13(59.1). 4 cases i.e18.9% cases were multiparous and 4(18.2) cases were nulliparous. the spectrum of presentation varied from pmb 9(40.9) cases, watery discharge 9(22) noscases, aub were 4(18.2) and pcb 4(18.2) TABLE1.

Of the total 22 cases all 22 (100)(%) underwent; laparotomy 20 (90.9) cases underwent type 2 radical hysterectomy,and 2 cases underwent type 3 radical hysterectomy. There were intraoperative surgical complications and 22(100) post operative complications. There wee 4(18.2) cases with a high residual urine and rest had minor bladdedydysfunction 18 (81.8) table -2.

The high residual urine post op for the four cases wee200 ml, 90ml, 150ml, 150ml. After adjuvant treatment all four case required re-catherterisation.

There were 2(9.09%) were adenosquamous ,1(4.55%) case was basaloid carcinoma and rest 20 (91%)cases were adenocarcinoma

The median size of the lesion is 3 cm .7 (33.3%) cases wee less

Table 1. Clinical Statistics.

Variables		Values
Age (in years)	Mean ± SD	46.04 ± 9.20
	Median	46
	Range (Max, Min)	34 (62, 28)
< 46, n (%)		09 (40.9)
≥ 46, n (%)		13 (59.1)
Parity, n (%)		
Nulliparity		04 (18.2)
Multiparity		18 (81.8)
Symptoms, n (%)		
PMB		09 (40.9)
WD		05 (22.7)
AUB		04 (18.2)
PCB		04 (18.2)

Table 2. Surgical Statistics.

Variables	Values
Surgical Procedure, n (%)	
Laparotomy	22 (100)
Radical Hysterectomy (type-2)	20 (90.9)
Radical Hysterectomy (type-3)	02 (09.1)
Laparoscopy	NIL
Surgical Complications, n (%)	
Intra op	NIL
Post op	22 (100)
Bowel	NIL
Bladder	22 (100)
Bladder Complications(HIGH RESIDUAL URINE)	04 (18.2)
Bladder Complications(MINOR BLADDER DYSFUNCTIONS)	18 (81.8)

Table 3. Histopathological Statistics.

Vacriables	Values	
Histology		
Adenosquamous	2	
Adenocarcinoma	20	
Basaloid carcinoma	1	
Size of lesion (in cm)	Mean ± SD	3.46 ± 1.5
	Median	3
	Range (Max, Min)	5.5 (7, 1.5)
< 3, n (%)	07 (33.3)	
≥ 3, n (%)	15 (66.7)	
Margins, n (%)		
Positive	10 (45.4)	
Negative	12 (54.6)	
LVSI, n (%)		
Positive	07 (33.3)	
Negative	15 (66.7)	
Grade, n (%)		
Grade-1	10 (45.4)	
Grade-2	06 (27.3)	
Grade-3	06 (27.3)	
Nodal Status, n (%)		
Pelvic		
Negative	14 (63.6)	
Positive	08 (36.4)	
Internal iliac	4	
Obturator	1	
Upper External iliac	2	
Low External iliac	1	
Common iliac	NIL	
Para aortic		
Negative	20 (90.9)	
Positive	02 (09.1)	
Depth of stomal invasion, n (%)		
< 5 mm	07 (31.8)	
≥5 mm	15 (68.2)	
Parametrium, n (%)		
Positive	05 (22.7)	
Negative	17 (77.3)	
Adnexa, n (%)		
Positive	NIL	
Negative	22 (100)	
Stage, n (%)		
Stage-1	17 (77.3)	
A	NIL	
B1	6	
B2	9	
B3	2	
Stage-2	NIL	
Stage-3	05 (22.7)	
C1	4	
C2	1	
Stage-4	NIL	

than 3 and 15 (66.7%) cases are more than 3 cm margins positive in 10 (45.4%) cases and negative in 12 (54.6%) lysi positive in 7 (31.8%) and 15 cases (66.7%) negative nos of grade 1 10(45.4) and grade 2 6(27.3%) and grade 3 6(27.3%)

Nodal status shows the pelvic node 8 (36.4%). Paraaortic 2(9.1%)

The depth of invasion >5mm 25(31.8%) and <5mm 7(68.2%)

The parametrium was positive in 5 cases (22.7%) cases was

negative in 17(77.3%) cases adnexa was not positive in any of cases

17 (77.3%) were in stage 1. With no case in 1a, 6 cases in 1b1, 9 cases in 1b2 and 2 cases in 1b3. There were 5 cases in stage 3. None of the cases presented in stage 2 and stage 4.

Survival Analysis

Figures

Figure 1. Adenocarcinoma of cervix.




Figure 2. The endocervical adenocarcinoma stained positive ihc pi6 +ve.




Table 4. Regression Analysis.

Coefficients:	Estimate (β)	Std. Error	z-value	OR	Pr(> z)
(Intercept)	-16.807	8.412	-1.998	0	0.046
Age	0.228	0.142	1.612	1.257	0.107
Pelvicnode	3.407	2.21	1.541	30.163	0.123
Sizeoflesion	0.778	0.512	1.519	2.177	0.129
Variable(s):					
Dependent Variable: Parametrium					
Predictor Variable: Age, Pelvicnode, Sizeoflesion.					

Not a single predictor variable is significant in predicting Parametrium Status.

Table 5. Regression Analysis5.

Coefficients:	Estimate (β)	Std. Error	z value	OR	Pr(> z)
(Intercept)	3.427	3.824	0.896	30.784	0.37
Age	-0.118	0.073	-1.607	0.889	0.108
Parametrium	2.637	1.775	1.486	13.97	0.137
Sizeoflesion	0.173	0.38	0.454	1.189	0.65
Depthofinvasion	-0.227	1.306	-0.174	0.797	0.862
Variable(s):					
Dependent Variable: Pelvicnode					
Predictor Variable: Age, Parametrium, Sizeoflesion, Depthofinvasion					

Not a single predictor variable is significant in predicting Pelvicnodal status.

Table 6. Survival Statistics.

Variables	Values
Lost To Follow-up, n (%)	03 (13.6)
NAD, n (%)	16 (72.8)
Death, n (%)	03 (13.6)

The nos of cases lost to follow up 3(13.6%).16 cases were normal on follow up and 3 cases succumbed (13.6%)

Table 7. Overall survival.

Time	No of Patients at risk	No of event(s)	Survival Probability	St. Error	lower 95% CI	upper 95% CI
48	15	1	0.933	0.064	0.815	1
50	12	1	0.856	0.095	0.688	1
55	10	1	0.77	0.118	0.57	1

Here, overall 5-year survival probability for the patients is 0.770 i.e 55 months with 95% CI (0.570, 1.000).

Table 7.1. Survival Status of the patients with Parametrium positive.

time	No of Patients at risk	No of event(s)	Survival Probability	St. Error	lower 95% CI	upper 95% CI
	48	12	1		0.917	0.079

Here, the 5-year survival probability for the patients with parametrium positive status is 0.917 48 months 95% CI (0.773, 1.000).

Table 8. Survival Status of the patients with Parametrium negative.


time	No of Patients at risk	No of event(s)	Survival Probability	St. Error	lower 95% CI	upper 95% CI
50	3	1	0.667	0.272	0.299	1
55	2	1	0.333	0.272	0.067	1

Here, the 5-year survival probability for the patients with parametrium negative status is 0.333i.e 55 months with 95% CI (0.067, 1.000).

Table 9. Log-rank test outcomes for survival differences.

Covariates	Category	Num. of Patient	Observed Event	Expect-ed Event	$(O-E)^2/E$	Chisq. Value	d.f.	p-value
Parametrium	Negative	18	1	2.35	0.776	3.59	1	0.06
	Positive	4	2	0.65	2.804			

The log rank test for change in survival between two parametrium groups i.e., negative and positive of cervical cancer patients gives chi-square value 3.59 (d.f.=1, p-value = 0.06), which indicates two groups do not differ in their survival.







Fig.2 Shows the proportion of individuals who have survived up to the specified time in the solid black line and their 95% confidence interval in black dashed lines.



Results

On regression analysis with dependent variable as the pelvic node was analysed with age, depth of invasion and size of lesion and parametrium, none of factors were significantly influencing the nodal status.

Similarly the parametrium positivity taken as the dependent vari-

able was analysed with age, size of lesion, depth of invasion. none of the factors could predict the parametrium involvement. overall 5-year survival probability for the patients is 0.770 i.e 55 months with 95% CI (0.570, 1.000). Here, the 5-year survival probability for the patients with parametrium positive status is 0.917 i.e 48 months 95% CI (0.773, 1.000). The log rank test for survival between two parametrium groups i.e., negative and positive patients gives chi-square value 3.59 (d.f.=1, p-value = 0.06), which indicates two groups do not differ in their survival.

Log-rank test outcomes for disease free survival differences

Covariates	Category	Num. of Patient	Observed Event	Expected Event	$\frac{(O - E)^2}{E}$	Chisq. Value	d.f.	p-value
parametrium	negative	18	1	2.21	0.662	2.8	1	0.1
	positive	4	2	0.79	1.851			

The log rank test for survival difference between two parametrium groups i.e., negative and positive of non squamous carcinoma patients gives chi-square value 2.8 (d.f.=1, p-value = 0.1), which indicates two groups do not differ in their disease free survival.

Disease free survival of the parametrium –ve/ parametrium +ve(60/24 mths). The log rank test for change in survival between two parametrium groups i.e., negative and positive patients gives chi-square value 2.8 (d.f.=1, p-value = 0.1), which indicates two groups do not differ in their disease free survival.

Conclusion

The parametrium and nodal positivity of non squamous cancers are not dependent on each other. Thus we conclude from overall survival of parametrium positive/parametrium negative (48mths/55mths) or the disease free survival of the two groups (24mths/60 mths) after radical hysterectomy with adjuvant crt. was not statistically significant. Although the parametrium negative cases showed increase in overall survival of 55mths and disease free survival of 60 mths.

Thus we can subject the patients to either of a single modal-

ity of treatment i.e either a radical hysterectomy or crt.

References

- [1]. Ali F, Kuelker R, Wassie B. Understanding cervical cancer in the context of developing countries. *Ann Trop Med Pub Health.* 2012 Jan 1;5(1).
- [2]. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer.* 2015 Mar 1;136(5):E359-86. PubMed PMID: 25220842.
- [3]. Bruni D, Angell HK, Galon J. The immune contexture and Immunoscoring in cancer prognosis and therapeutic efficacy. *Nature Reviews Cancer.* 2020 Nov;20(11):662-80.
- [4]. Bellavite P, Ortolani R, Pontarollo F, Piasere V, Benato G, Conforti A. Immunology and homeopathy. 4. Clinical studies-part 2. *Evidence-Based Complementary and Alternative Medicine.* 2006 Dec 1;3(4):397-409.
- [5]. eifeljp et al. 1990.
- [6]. Nasri MN, Setchell ME, Chard T. Transvaginal ultrasound for diagnosis of uterine malformations. *Br J ObstetGynaecol.* 1990 Nov;97(11):1043-5. PubMed PMID: 2252870.
- [7]. Mabuchi et al. *International Journal of Oncology.* 2001.
- [8]. Gloria salvo. nectur study ijc. 2021.